

Dietary Reference Intakes: Elements

Nutrient	Function	Life Stage Group	RDA/AI*	UL ^a	Selected Food Sources	Adverse effects of excessive consumption	Special Considerations
Calcium	Essential role in blood clotting, muscle contraction, nerve transmission, and bone and tooth formation	Infants	(mg/d)	(mg/d)	Milk, cheese, yogurt, corn tortillas, calcium-set tofu, Chinese cabbage, kale, broccoli	Kidney stones, hypercalcemia, milk alkali syndrome, and renal insufficiency	Amenorrheic women (exercise- or anorexia nervosa-induced) have reduced net calcium absorption. There is no consistent data to support that a high protein intake increases calcium requirement.
		0–6 mo	210*	ND ^b			
		7–12 mo	270*	ND			
		Children					
		1–3 y	500*	2,500			
		4–8 y	800*	2,500			
		Males					
		9–13 y	1,300*	2,500			
		14–18 y	1,300*	2,500			
		19–30 y	1,000*	2,500			
		31–50 y	1,000*	2,500			
		50–70 y	1,200*	2,500			
		> 70 y	1,200*	2,500			
		Females					
		9–13 y	1,300*	2,500			
		14–18 y	1,300*	2,500			
		19–30 y	1,000*	2,500			
		31–50 y	1,000*	2,500			
		50–70 y	1,200*	2,500			
		> 70 y	1,200*	2,500			
Pregnancy							
≤ 18 y	1,300*	2,500					
19–30y	1,000*	2,500					
31–50 y	1,000*	2,500					
Lactation							
≤ 18 y	1,300*	2,500					
19–30y	1,000*	2,500					
31–50 y	1,000*	2,500					
Chromium	Helps to maintain normal blood glucose levels	Infants	(µg/d)		Some cereals, meats, poultry, fish, beer	Chronic renal failure	None
		0–6 mo	0.2*	ND			
		7–12 mo	5.5*	ND			
		Children					
		1–3 y	11*	ND			
		4–8 y	15*	ND			
		Males					
		9–13 y	25*	ND			
		14–18 y	35*	ND			
		19–30 y	35*	ND			
		31–50 y	35*	ND			
		50–70 y	30*	ND			
		> 70 y	30*	ND			
		Females					
		9–13 y	21*	ND			
		14–18 y	24*	ND			
		19–30 y	25*	ND			
		31–50 y	25*	ND			
		50–70 y	20*	ND			
		> 70 y	20*	ND			
Pregnancy							
≤ 18 y	29*	ND					
19–30y	30*	ND					
31–50 y	30*	ND					
Lactation							
≤ 18 y	44*	ND					
19–30y	45*	ND					
31–50 y	45*	ND					

NOTE: The table is adapted from the DRI reports, see www.nap.edu. It represents Recommended Dietary Allowances (RDAs) in **bold type**, Adequate Intakes (AIs) in ordinary type followed by an asterisk (*), and Tolerable Upper Intake Levels (ULs)^a. RDAs and AIs may both be used as goals for individual intake. RDAs are set to meet the needs of almost all (97 to 98 percent) individuals in a group. For healthy breastfed infants, the AI is the mean intake. The AI for other life stage and gender groups is believed to cover the needs of all individuals in the group, but lack of data prevent being able to specify with confidence the percentage of individuals covered by this intake.

^aUL = The maximum level of daily nutrient intake that is likely to pose no risk of adverse effects. Unless otherwise specified, the UL represents total intake from food, water, and supplements. Due to lack of suitable data, ULs could not be established for vitamin K, thiamin, riboflavin, vitamin B₁₂, pantothenic acid, biotin, or carotenoids. In the absence of ULs, extra caution may be warranted in consuming levels above recommended intakes.

^bND = Not determinable due to lack of data of adverse effects in this age group and concern with regard to lack of ability to handle excess amounts. Source of intake should be from food only to prevent high levels of intake.

SOURCES: *Dietary Reference Intakes for Calcium, Phosphorous, Magnesium, Vitamin D, and Fluoride* (1997); *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B₆, Folate, Vitamin B₁₂, Pantothenic Acid, Biotin, and Choline* (1998); *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids* (2000); and *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc* (2001). These reports may be accessed via www.nap.edu. Copyright 2001 by the National Academies. All rights reserved.

